

AGRIMEC GOLD

8.4 % SC

**lowest dose...
Get rid of the dust mites**



Active ingredient :-

(Abamectin 8.4% SC)

Introduction :-

A broad-spectrum insecticide / acaricide for mites (motile stages only). It acts as an infectious poison and has a lesser effect on contact. It also has the ability to penetrate the leaves of plants.. It activates the action of gamma-amino acid, which leads to preventing the transmission of nerve signals, and then the insects stop and The dream is about movement and activity, and thus stopping feeding and laying eggs, and it dies.. and does not cause any damage to the plant until it is eliminated.

Photos of injuries on different crops



Dust mites in date palm



Thrips




Leaf miner



Mites in tomato



Mites in cucumber




Mites in pepper



Mites in ornamental plants

Advantages of using Agrimec Gold :-

- 1) It contains an active substance produced in nature for some organisms that live in the soil .
- 2) It quickly penetrates the epidermis of the plant. It quickly permeates the surface of the leaves and is stored inside .
- 3) It has a short residual effect on your crop, which leads to the production of a high quality crop .
- 4) Agrimec Gold 4.8% SC - as an active substance - is characterized by having 8 isomers, only two are active and the rest are either ineffective or have weak efficacy. Agrimec Gold 4.8% SC contains only the two active isotopes of abamectin, compared to other sources of abamectin, which have a mixture of isotopes, which makes the effectiveness of Agrimec Gold 4.8% high compared to other sources .



Usage Recommendations :-

- Safe on natural, vital enemies... to combat mites, spiders and tunnel makers
- Spraying is done at the onset of infection.. The spraying pressure and the solution must be sufficient to cover all parts of the plant, including the lower surface of the leaf, well with spraying.
- To ensure long-term effectiveness, the pesticide is mixed with a spreading agent
- Do not spray more than 2 sprays for grapes and apples - 3 sprays for citrus fruits in one cycle.
- It is not recommended to use the substance on cloves (ornamental plant).
- It is not allowed to graze livestock in treated areas or provide treated plants before 28 days have passed since spraying.

Dosage :-

Crop	Target pests	Dose	PHI
Vegetables (Eggplant-tomato-sweet pepper-potato-pumpkins)	Liriomyza SPP (Leafminers)	15ml / 100 L	3
Cucumber	Tetranychus urticae	15ml / 100 L	7
Strawberry	Spider mite	150ml / Ha (Not less than 500 L/ Ha)	7
Fruit trees	Phyllocoptruta Oleivora	12.5ml / 100 L	10
pear	Eotteranychus spp (Yellow mite)	150 - 200ml / 1000 : 2000 L / Ha	10
	Spider mite		
Cut-flowers and Orna- mental plants	(Leaf miner)	12.5ml / 100 L	-
	Spider mite	15ml / 100 L	

Packing :-

Agrimec Gold is available in 100 ml